

Upper airway disorders within adult respiratory services – guidance

Key points

- The term upper airway disorders within adult respiratory services is used to differentiate between upper and lower respiratory disorder and broader speech and language therapy clinical populations related to dysphagia
- SLTs are specialists in the assessment, diagnosis and treatment of upper airway disorders, which includes inducible laryngeal obstruction and chronic cough
- Speech and language therapy for patients with upper airway disorders improves patient-reported quality of life and reduces healthcare use significantly; patients should have equitable access to appropriately trained SLTs.
- SLTs working in upper airway disorders are members of a specialised respiratory multidisciplinary team, and it is recommended they should be commissioned as such.
- Dysphonia, in the absence of any respiratory symptom burden, should be managed according to the [**RCSLT voice guidance**](#) and within the context of a multidisciplinary team (MDT) ENT setting.

Please also read RCSLT position paper: [**The role of speech and language therapy in upper airway disorders within adult respiratory services \(PDF\)**](#).

Clinical population

Upper airway disorders represent abnormal laryngeal functions, which are thought to develop as a result of heightened laryngeal sensitivity and hyper-responsiveness (Famokunwa et al, 2019; Hull et al, 2016; Sundar et al, 2021). They include:

- inducible laryngeal obstruction (ILO)
- chronic cough (CC)
- persistent throat clearing
- globus pharyngeus
- heightened laryngopharyngeal sensitivity symptoms (eg throat irritation, atypical throat sensations)
- dysphagia symptoms (without evidence of impaired swallow function)
- dysphonia

When patterns of abnormal laryngeal function occur a number of respiratory symptoms can develop (eg cough, breathlessness, wheeze). The umbrella term 'laryngeal dysfunction' captures these symptoms, which often appear refractory to treatment and incongruous with a clinical disease state (Hull et al, 2016).

Patients often present with an overlap of upper airway disorder symptoms (Vertigan et al, 2013), which suggests the manifestations of laryngeal dysfunction may be mechanistically linked (Sundar et al, 2021). Evidence supports this theory as the clinical profiles of ILO and CC are markedly similar in terms of demographic and clinical attributes (Ryan et al, 2009).

Inducible laryngeal obstruction

Inducible laryngeal obstruction describes inappropriate transient laryngeal closure during respiration. In the absence of any structural or neurological abnormalities, airflow obstruction occurring at the glottic and/or supraglottic level leads to breathing difficulties (Halvorsen et al, 2017). Understanding of the condition remains limited as available evidence is largely retrospective. Patients present across various healthcare settings with differing levels of morbidity, ranging from mild dyspnoea to acute respiratory distress (Haines et al, 2018).

Symptoms are sudden in onset, often triggered by exposure to an external environmental trigger, and typically affect inspiration (with associated throat tightness and often noisy breathing) (Dunn et al, 2015; Halvorsen et al, 2017). Due to commonality in symptoms ILO is often misdiagnosed as asthma (Haines et al, 2018; Newman et al, 1994; Newman et al, 1995). To add to the complexity the two conditions are not mutually exclusive and can co-occur (Low et al, 2011; McDonald, 2019).

Historically, numerous terms (including vocal cord dysfunction and paradoxical vocal fold motion/disorder) have been applied to describe inappropriate laryngeal closure (Christensen et al, 2015; Haines et al, 2018), which has caused significant confusion and hindered research developments. The European Respiratory Society, European Laryngological Society, and the American College of Chest Physicians led an international task force in 2015 with an aim to standardise nomenclature (Christensen et al, 2015). International consensus was achieved, and the term 'inducible laryngeal obstruction' proposed with clear justification and explanation.

The RCSLT supports the adoption of this taxonomy and recommends SLTs working within the field of upper airway disorders use inducible laryngeal obstruction (ILO) routinely.

Chronic cough

Cough is a protective reflex mechanism, which enables airway secretion clearance and prevents aspiration. The majority of cases of cough are acute or subacute and arise due to viral upper respiratory tract infections, usually lasting less than three weeks. However, when a cough persists for more than eight weeks it is defined as 'chronic cough' (Irwin et al, 2006).

Chronic cough patients classically describe a dry irritable cough, which triggers in response to environmental irritants and a subsequent urge to cough (Hilton et al, 2013). Due to research advances, it is now accepted CC represents a hyperresponsiveness of the neuronal pathways involved in the cough reflex (Satia et al, 2017), and an impairment in descending inhibitory controls (Ando et al, 2016; Farrell et al, 2012). More recently the phenomena have been referred to as cough hypersensitivity syndrome (Morice et al, 2014).

A systematic approach to diagnosis and treatment can be successful but cough can remain refractory in approximately 20% of cases (Pratter and Abouzgheib, 2006). Persistent cough causes significant physical, psychological and social morbidity. Many patients suffer from incontinence, vomiting and depression, thus negatively affecting quality of life (French et al, 1998).

Role of speech and language therapy

The remit of SLTs working in upper airway disorders within respiratory services is both broad and specialised. SLTs are required to manage patients with multiple and complex comorbidities and work across dedicated and disparate MDT teams. The speech and language therapy workforce has the skills to contribute across a patient's clinical journey; ie assessment, differential diagnosis, management planning, treatment and discharge.

SLTs play a key role in educating patients to be able to differentiate respiratory symptoms and therefore maximise effectiveness of medical interventions. Speech and language therapy is frequently referred to as the 'gold standard' beneficial treatment for ILO (Kenn and Balkissoon, 2011; Marcinow et al, 2015; Patel et al, 2015), despite the lack of prospective randomised control trials. In CC, better quality evidence exists and supports the role of SLTs in managing CC patients; in two randomised control trials speech and language therapy reduced cough frequency and improved health-related quality of life (Chamberlain et al, 2017; Vertigan et al, 2006).

Workforce planning within organisations should occur to ensure SLTs are available to provide high-quality care for patients with upper airway disorders. SLTs working in upper airway disorders are members of a specialised respiratory MDT, and it is recommended they should be commissioned as such.

Impact of upper airway disorders and benefit of speech and language therapy

Upper airway disorders are associated with high healthcare utilisation, which negatively impacts healthcare resources and patient morbidity. No robust data exists to quantify the economic impact, but the burden is likely significant. Speech and language therapy can improve outcomes on patient-recorded quality of life (Bassil et al, 2018; Chamberlain et al, 2017; Fowler et al, 2015; Patel et al, 2015; Pargeter and Mansur, 2016; Slinger et al, 2019; Vertigan et al, 2006) and significantly reduce healthcare usage (Murphy et al, 2020). Therefore, the need for speech and language therapy provision to support alleviation of symptoms is self-explanatory.

Inducible laryngeal obstruction

Data from the UK national ILO registry (Haines et al, 2020) identifies high healthcare utilisation within the population; 69% of individuals had visited emergency departments at least once in 12 months, with an average of five attendances. Critical care admission occurred in one in five patients, with 10% receiving intubation and ventilatory support. Patient morbidity was significant; over half were off work on long-term sickness due to their symptoms and 64% reported impaired functional capacity on the MRC breathlessness scale.

According to a recent systematic review, patients with asthma and co-existing ILO demonstrate greater healthcare use by more than a third, compared to those with asthma only (Murphy et al, 2020). Nearly 40% of patients suffering with ILO were misdiagnosed as asthmatic for an average 5.3 years, resulting in inappropriate pharmacological burden.

Chronic cough

Chronic cough affects between 5-10% of the adult population (Song et al, 2015), and is responsible for up to 40% of respiratory outpatient referrals (Pratter et al, 2006). The impact of CC on a patient is significant and includes physical complications (eg vomiting, incontinence, cracked ribs) as well as psychosocial problems (eg social isolation). When compared with other chronic medical conditions, CC sufferers have significantly higher levels of depression due to symptom burden (Adams et al, 2009).

Multidisciplinary working

Multidisciplinary team working is an integral role for SLTs working in upper airways disorders. During differential diagnosis, SLTs inform the MDT of the contribution upper airway function is playing on respiration and suggest appropriate strategies to maximise function when required. Regular input to specialist airway MDT meetings is a common role. Core MDT colleagues include:

- SLTs
- respiratory physicians
- respiratory physiotherapists
- respiratory clinical nurse specialists
- lung physiologist
- mental health practitioners/clinical psychologists
- otolaryngologists

Wider MDT colleagues include:

- allergists
- cardiothoracic physicians
- critical care intensivists
- dietitians
- gastroenterologists
- oesophageal laboratory technicians
- pharmacists

SLTs working in upper airway disorders are often members of a specialised respiratory MDTs, and it is recommended they should be commissioned as such.

Referral

Cough, breathlessness and other respiratory symptoms can arise as a symptom of a number of conditions (eg malignancy, airway stenosis, uncontrolled asthma, bronchiectasis). It is therefore imperative, prior to any speech and language therapy intervention, that thorough assessment from a respiratory physician occurs. This assessment may take place in a uni-professional respiratory clinic or as part of a dedicated upper airways MDT respiratory clinic. Referrals containing any red flag symptoms (eg haemoptysis, unintentional weight loss, persistent noisy breathing) should be managed in accordance with existing professional guidelines and re-directed onto relevant pathways.

Referral pathways will vary according to local services and commissioning. In specialist centres, initial referrals may be for MDT respiratory assessment rather than isolated speech and language therapy assessment, with subsequent referral for ongoing speech and language therapy management as a result of the MDT assessment process. SLTs may receive direct referrals from respiratory clinics once respiratory assessment has occurred (especially in CC populations). However, in all cases, referrals should provide as much existing data as possible on previous investigations to support assessment and ongoing management.

For SLTs working in upper airway disorders, the minimum recommended referral criteria include:

- respiratory medical review within the last six months
- ability to refer directly back for respiratory review (with a low threshold applied), if no longer under the care of a dedicated respiratory physician
- endoscopic evaluation of the larynx (continuous laryngoscopy in suspected ILO cases) within the last six months
- optimised management of associated conditions (eg asthma)
- patient being aware of diagnosis and consents to referral
- patient being appropriate for speech and language intervention and is likely to benefit
- patient is aware of the role of speech and language therapy in managing upper airway disorders and is happy to engage

Based on the above minimum criteria, SLTs are unlikely to regularly accept direct referrals from GPs (eg direct GP referral for CC management would be rejected if there had been no respiratory review in the previous six months). Similarly, self-referrals should be rejected.

Assessment

Clinical evaluation

It is important to gain a comprehensive understanding of previous events, current status and patient expectations. The minimum case history data set should include:

- presenting complaint
- history of presenting complaint
- significant past medical history and diagnosed co-morbidities (eg reflux, nasal disease)
- drug history (particularly use of asthma medications and angiotensin-converting enzyme inhibitors)
- occupational history
- social history including smoking and employment status
- respiratory specific details (eg symptom onset, triggers – including any environmental, exacerbations, mucous burden, treatment trials and responses, previous intubations)

Investigations

Investigations will be tailored to the patient and directed by respiratory physicians. As a minimum, prior to speech and language therapy intervention, investigation should include:

- endoscopic evaluation of the larynx (and continuous laryngoscopy in suspected inducible laryngeal cases)
- imaging (chest x-ray)
- lung physiology (spirometry, \pm reversibility, flow volume loops)

Other assessment investigations may include:

- breathing pattern disorder assessment
- bronchoscopy
- cardiac specific investigations (eg electrocardiogram)
- data related to allergy (immunoglobulin E, skin prick test) and inflammation (blood/sputum eosinophils, fractional exhaled nitric oxide)
- gastroenterology specific investigations (eg oesophageal manometry and 24-hour ph monitoring/barium swallow)
- imaging (high-resolution computed tomography)
- objective dysphagia assessments

Treatment

Speech and language therapy interventions are typically multi-modal and may be combined with other treatments (eg joint working with respiratory physiotherapists when there is a co-existing breathing pattern disorder). Typically, speech and language therapy will be delivered via a one-to-one method to facilitate a high level of personalisation. However, group therapy for the management of CC has been reported effective (Selby et al, 2018).

Generally, strategies aim to facilitate reduction in laryngeal irritants/aggravating behaviours, behavioural modifications and distraction techniques. Specifically in ILO the primary role is to teach control of the laryngeal area and maintain an adequately open airway during respiration. In CC, there is usually a greater emphasis on cough suppression techniques.

Common components used in speech and language therapy management of upper airway disorders include (not an exhaustive list):

- cough suppression
- diaphragmatic breathing
- differentiation of symptoms
- discussion of contributory and maintaining factors
- education and explanation
- laryngeal airway control exercises (including emergency 'release' breathing techniques)
- psychoeducational counselling (including goal setting)
- relaxation of head and neck extrinsic muscle tension
- release of inappropriate upper airway constriction
- supporting decision making for pharmacological interventions, as part of an MDT (eg biological therapy for severe asthma, reduction in inappropriate steroid use)
- upper airway health
- visual-biofeedback therapeutic laryngoscopy
- Controlling symptoms during purposeful trigger exposure

Monitoring

Several symptom questionnaires exist to evaluate upper airway disorders. Key questionnaires include:

- Leicester Cough Questionnaire (Birring et al, 2003)
- Cough Severity Index (Shambel et al, 2013)
- VCDQ (Fowler et al, 2015)
- Newcastle laryngeal hypersensitivity questionnaire (Vertigan et al, 2014)
- Pittsburgh vocal cord dysfunction index (Traister et al, 2014)
- Dyspnoea index (Witek et al, 2003)

Supplementary questionnaires may include (not an exhaustive list):

- Reflux symptom index (Belafsky et al, 2002)
- Nijmegen Questionnaire (Van Doorn et al, 1982)
- Brompton Breathing Pattern Assessment Tool (Todd et al, 2018)
- EAT-10 (Belafsky et al, 2008)
- Voice Handicap Index (Rosen et al, 2004)
- Voice Symptom Score (Wilson et al, 2004)

The RCSLT has adopted **Therapy Outcome Measures** (TOMS) (Enderby et al, 2013) as the profession's outcome measurement tool. To support SLTs working in upper airway disorders, a group of senior SLTs are developing a standardised TOMS framework for upper airway disorders. The final validation of the tool is underway and will be available here in due course.

In line with the requirements of the Health and Care Professions Council (HCPC), ongoing monitoring should occur. This will facilitate an understanding of progress and provide data on the effectiveness of speech and language therapy intervention to support ongoing management and appropriate discharge (together with patient therapy goals).

For patients with known co-existing respiratory disease it is essential that if treatment response is not as expected, there is deterioration in respiratory symptoms or there are any other concerns, respiratory review should occur prior to continued speech and language therapy intervention, preferably within the MDT context.

Continuous laryngoscopy

Endoscopic evaluation of the larynx enables endoscopic evaluation of the nasal cavities, pharyngeal space and laryngeal vestibule. It establishes if there are any structural, pathological, neurological or functional abnormalities. However, in the context of ILO, it only provides a snapshot in time and in the absence of concurrent respiratory symptoms, the laryngopharynx functions normally.

The frequently cited gold standard for ILO diagnosis is laryngoscopy during a symptomatic episode (Halvorsen et al, 2017). However, diagnostic confirmation is dependent on being able to perform laryngoscopy during a symptomatic attack. This is not always logistically possible and highly dependent on direct accessibility of equipment and appropriately trained clinicians. Therefore, in recent years provocation agents have been used on individuals who are baseline asymptomatic, with an aim to induce and replicate compatible clinical features within a controlled environment.

Continuous laryngoscopy during provocation (Hull et al, 2019) or exercise (Heimdal et al, 2006) provides a more targeted assessment as continuous monitoring occurs while symptom inducers are presented. It enables visualisation of any laryngeal reactions within the laryngeal vestibule in real time. Diagnostic yields in this context are reported to be more accurate and targeted (Halvorsen et al, 2017; Hull et al, 2019). However, despite efforts to provoke symptoms during continuous laryngoscopy, individuals may remain asymptomatic. It is therefore important to ensure false negative ILO diagnoses are not given; the confirmation of an ILO diagnosis during continuous laryngoscopy is only possible when the individual is symptomatic. When no symptoms are elicited during the procedure, the MDT should look to the presence or absence of other clinical signs to inform differential diagnosis (eg patient history/description of symptoms/truncation of inspiratory flow loops).

The **RCSLT upper airways position paper (PDF)** provides the professional clinical context within which SLTs should practice continuous laryngoscopy and is the RCSLT's official statement of professional practice for SLTs using it. It is the professional responsibility of individual SLTs to ensure adherence to its content; evidence of such adherence will help to ensure professional indemnity. Guidance includes scope of practice, purpose, procedure, patient information, consent, equipment, assessment protocols and image interpretation guidance.

References

- Adams RJ, Appleton SL, Wilson DH, Taylor AW, Ruffin RE. Associations of physical and mental health problems with chronic cough in a representative population cohort. *Cough*. 2009 Dec;5(1):1-9.
- Ando A, Smallwood D, McMahon M, Irving L, Mazzone SB, Farrell MJ. Neural correlates of cough hypersensitivity in humans: evidence for central sensitisation and dysfunctional inhibitory control. *Thorax*. 2016 Apr 1;71(4):323-9.
- Bassil L, Pargeter N, Fellows J, Howard R, Mansur A. Quality of life in inducible laryngeal obstruction; a questionnaire pilot. *Thorax Supl*. 2018 73 (4:M30)
- Belafsky PC, Mouadeb DA, Rees CJ, Pryor JC, Postma GN, Allen J, Leonard RJ. Validity and reliability of the Eating Assessment Tool (EAT-10). *Annals of Otology, Rhinology and Laryngology*. 2008 Dec;117(12):919-24.
- Belafsky PC, Postma GN, Koufman JA. Validity and reliability of the reflux symptom index (RSI). *Journal of voice*. 2002 Jun 1;16(2):274-7.
- Birring SS, Prudon B, Carr AJ, Singh SJ, Morgan MD, Pavord ID. Development of a symptom specific health status measure for patients with chronic cough: Leicester Cough Questionnaire (LCQ). *Thorax*. 2003 Apr 1;58(4):339-43.
- Chamberlain MS, Garrod R, Clark L, Douiri A, Parker SM, Ellis J, Fowler SJ, Ludlow S, Hull JH, Chung KF, Lee KK. Physiotherapy, and speech and language therapy intervention for patients with refractory chronic cough: a multicentre randomised control trial. *Thorax*. 2017 Feb;72(2):129.
- Christensen PM, Heimdahl JH, Christopher KL, Bucca C, Cantarella G, Friedrich G, Halvorsen T, Herth F, Jung H, Morris MJ, Remale M. ERS/ELS/ACCP 2013 international consensus conference nomenclature on inducible laryngeal obstructions. *European Respiratory Review*. 2015 Sep 1;24(137):445-50.
- Dunn NM, Katial RK, Hoyte FC. Vocal cord dysfunction: a review. *Asthma research and practice*. 2015 Dec;1(1):1-8.
- Enderby P, John A, Petheram B. Therapy outcome measures for rehabilitation professionals: speech and language therapy, physiotherapy, occupational therapy. John Wiley & Sons; 2013 May 31.
- Famokunwa B, Walsted ES, Hull JH. Assessing laryngeal function and hypersensitivity. *Pulmonary pharmacology and therapeutics*. 2019 Jun 1;56:108-15.
- Farrell MJ, Cole LJ, Chiapoco D, Egan GF, Mazzone SB. Neural correlates coding stimulus level and perception of capsaicin-evoked urge-to-cough in humans. *Neuroimage*. 2012 Jul 16;61(4):1324-35.

Fife TA, Butler SG, Langmore SE, Lester S, Wright Jr SC, Kemp S, Grace-Martin K, Rees Lintzenich C. Use of topical nasal anesthesia during flexible endoscopic evaluation of swallowing in dysphagic patients. *Annals of Otolaryngology, Rhinology and Laryngology*. 2015 Mar;124(3):206-11.

Forrest LA, Husein T, Husein O. Paradoxical vocal cord motion: classification and treatment. *The Laryngoscope*. 2012 Apr;122(4):844-53.

Fowler SJ, Thurston A, Chesworth B, Cheng V, Constantinou P, Vyas A, Lillie S, Haines J. The VCDQ—a Questionnaire for symptom monitoring in vocal cord dysfunction. *Clinical & Experimental Allergy*. 2015 Sep;45(9):1406-11.

French CT, Irwin RS, Fletcher KE, Adams TM. Evaluation of a cough-specific quality-of-life questionnaire. *Chest*. 2002 Apr 1;121(4):1123-31.

Gibson PG, Chang AB, Glasgow NJ, Holmes PW, Kemp AS, Katelaris P, Landau LI, Mazzone S, Newcombe P, Van Asperen P, Vertigan AE. CICADA: Cough in Children and Adults: Diagnosis and Assessment. Australian cough guidelines summary statement. *Medical Journal of Australia*. 2010;192(5):265-71.

H, Zorko R. Fibreoptic Endoscopic evaluation of Swallowing (FEES)

Haines J, Esposito K, Slinger C, Pargeter N, Murphy J, Selby J, Prior K, Mansur A, Vyas A, Stanton AE, Sabroe I. UK consensus statement on the diagnosis of inducible laryngeal obstruction in light of the COVID 19 pandemic. *Clinical & Experimental Allergy*. 2020 Dec;50(12):1287-93.

Haines J, Hull JH, Fowler SJ. Clinical presentation, assessment, and management of inducible laryngeal obstruction. *Current opinion in otolaryngology & head and neck surgery*. 2018 Jun 1;26(3):174-9.

Haines J, Fowler SJ, Simpson AJ, Selby J, Hull JH. High healthcare utilisation and functional impairment in inducible laryngeal obstruction: results from the UK national registry. 2020. ERS abstract presentation.

Halvorsen T, Walsted ES, Bucca C, Bush A, Cantarella G, Friedrich G, Herth FJ, Hull JH, Jung H, Maat R, Nordang L. Inducible laryngeal obstruction: an official joint European Respiratory Society and European Laryngological Society statement. *European Respiratory Journal*. 2017 Sep 1;50(3).

Heimdal JH, Roksund OD, Halvorsen T, Skadberg BT, Olofsson J. Continuous laryngoscopy exercise test: a method for visualizing laryngeal dysfunction during exercise. *The Laryngoscope*. 2006 Jan;116(1):52-7.

Hilton EC, Baverel PG, Woodcock A, Van Der Graaf PH, Smith JA. Pharmacodynamic modeling of cough responses to capsaicin inhalation calls into question the utility of the C5 end point. *Journal of allergy and clinical immunology*. 2013 Oct 1;132(4):847-55.

Hull JH, Backer V, Gibson PG, Fowler SJ. Laryngeal dysfunction: assessment and management for the clinician. *American journal of respiratory and critical care medicine*. 2016 Nov 1;194(9):1062-72.

Hull JH, Walsted ES, Feary J, Cullinan P, Scadding G, Bailey E, Selby J. Continuous laryngoscopy during provocation in the assessment of inducible laryngeal obstruction. *The Laryngoscope*. 2019 Aug;129(8):1863-6.

Hull JH. Multidisciplinary team working for vocal cord dysfunction: Now it's GO time. 2019 714-715

Irwin RS, Baumann MH, Bolser DC, Boulet LP, Braman SS, Brightling CE, Brown KK, Canning BJ, Chang AB, Dicipinigaitis PV, Eccles R. Diagnosis and management of cough executive summary: ACCP evidence-based clinical practice guidelines. *Chest*. 2006 Jan 1;129(1):1S-23S.

Irwin RS, French CT, Lewis SZ, Diekemper RL, Gold PM, Adams TM, Altman KW, Barker AF, Birring SS, Bolser DC, Boulet LP. Overview of the management of cough: CHEST Guideline and Expert Panel Report. *Chest*. 2014 Oct 1;146(4):885-9.

Jones SM, Awad R, Esposito K, Shaw J, Slade S, Stewart C, Young K. Speech and Language Therapy Endoscopic Evaluation of the Larynx for Clinical Voice Disorders. London: Royal College of Speech and Language Therapists, position paper, 2020.

Jones SM, Awad R, Esposito K, Shaw J, Slade S, Stewart C, Young K. Speech and Language Therapy Endoscopic Evaluation of the Larynx for Clinical Voice Disorders. London: Royal College of Speech and Language Therapists, competency framework and training log, 2020.

Kenn K, Balkissoon R. Vocal cord dysfunction: what do we know?. *European Respiratory Journal*. 2011 Jan 1;37(1):194-200.

Lin J, Walsted ES, Backer V, Hull JH, Elson DS. Quantification and analysis of laryngeal closure from endoscopic videos. *IEEE Transactions on Biomedical Engineering*. 2018 Aug 29;66(4):1127-36.

Low K, Lau KK, Holmes P, Crossett M, Vallance N, Phyland D, Hamza K, Hamilton G, Bardin PG. Abnormal vocal cord function in difficult-to-treat asthma. *American journal of respiratory and critical care medicine*. 2011 Jul 1;184(1):50-6.

Marcinow AM, Thompson J, Forrest LA, deSilva BW. Irritant-induced paradoxical vocal fold motion disorder: diagnosis and management. *Otolaryngology-Head and Neck Surgery*. 2015 Dec;153(6):996-1000.

McDonald VM, Hiles SA, Godbout K, Harvey ES, Marks GB, Hew M, Peters M, Bardin PG, Reynolds PN, Upham JW, Baraket M. Treatable traits can be identified in a severe asthma registry and predict future exacerbations. *Respirology*. 2019 Jan;24(1):37-47.

Morice AH, Millqvist E, Belvisi MG, Bieksiene K, Birring SS, Chung KF, Dal Negro RW, Dicpinigaitis P, Kantar A, McGarvey LP, Pacheco A. Expert opinion on the cough hypersensitivity syndrome in respiratory medicine. *European Respiratory Journal*. 2014 Nov 1;44(5):1132-48.

Morice AH, Millqvist E, Bieksiene K, Birring SS, Dicpinigaitis P, Ribas CD, Boon MH, Kantar A, Lai K, McGarvey L, Rigau D. ERS guidelines on the diagnosis and treatment of chronic cough in adults and children. *European Respiratory Journal*. 2020 Jan 1;55(1).

Murphy JM, Stephen S, Pearson F, DeSoyza A. P108 A systematic review to explore the relationship between inducible laryngeal obstruction and healthcare utilisation in adults with Asthma.

Newman KB, Dubester SN. Vocal cord dysfunction: masquerader of asthma. In *Seminars in Respiratory and Critical Care Medicine* 1994 Mar (Vol. 15, No. 02, pp. 161-167). Copyright© 1994 by Thieme Medical Publishers, Inc..

Newman KB, Mason 3rd UG, Schmalig KB. Clinical features of vocal cord dysfunction. *American journal of respiratory and critical care medicine*. 1995 Oct;152(4):1382-6.

Pargeter N, Mansur AH. P226 Vocal cord dysfunction; clinical outcomes of speech and language therapy intervention. *Thorax Supl*. 2016 71 (3: P226) .

Patel RR, Venediktov R, Schooling T, Wang B. Evidence-based systematic review: effects of speech-language pathology treatment for individuals with paradoxical vocal fold motion. *American journal of speech-language pathology*. 2015 Aug;24(3):566-84.

Pratter MR, Abouzgheib W. Make the cough go away. *Chest*. 2006 May 1;129(5):1121-3.

RCSLT (2019). Dsdas. **[Giving Voice to people with upper airway disorders \(PDF\)](#)**.

Rosen CA, Lee AS, Osborne J, Zullo T, Murry T. Development and validation of the voice handicap index 10. *The Laryngoscope*. 2004 Sep;114(9):1549-56.

Ryan NM, Gibson PG. Characterization of laryngeal dysfunction in chronic persistent cough. *The Laryngoscope*. 2009 Apr;119(4):640-5.

Satia I, Tsamandouras N, Holt K, Badri H, Woodhead M, Ogungbenro K, Felton TW, O'Byrne PM, Fowler SJ, Smith JA. Capsaicin-evoked cough responses in asthmatic patients: evidence for airway neuronal dysfunction. *Journal of Allergy and Clinical Immunology*. 2017 Mar 1;139(3):771-9.

Selby J, Hull JH, Bailey E, Tidmarsh B. P4 A new cough remedy? patient evaluation of cough therapy group intervention. 2018 Thorax suppl A97-A98

Shembel AC, Rosen CA, Zullo TG, Gartner Schmidt JL. Development and validation of the cough severity index: a severity index for chronic cough related to the upper airway. The Laryngoscope. 2013 Aug;123(8):1931-6.

Slinger C, Mehdi SB, Milan SJ, Dodd S, Matthews J, Vyas A, Marsden PA. Speech and language therapy for management of chronic cough. Cochrane Database of Systematic Reviews. 2019(7).

Smith JA, Kitt MM, Morice AH, Birring SS, McGarvey LP, Sher MR, Li YP, Wu WC, Xu ZJ, Muccino DR, Ford AP. Gefapixant, a P2X3 receptor antagonist, for the treatment of refractory or unexplained chronic cough: a randomised, double-blind, controlled, parallel-group, phase 2b trial. The Lancet Respiratory Medicine. 2020 Aug 1;8(8):775-85.

Song WJ, Chang YS, Faruqi S, Kim JY, Kang MG, Kim S, Jo EJ, Kim MH, Plevkova J, Park HW, Cho SH. The global epidemiology of chronic cough in adults: a systematic review and meta-analysis. European Respiratory Journal. 2015 May 1;45(5):1479-81.

Sundar KM, Stark AC, Hu N, Barkmeier-Kraemer J. Is laryngeal hypersensitivity the basis of unexplained or refractory chronic cough?. ERJ Open Research. 2021 Jan 1;7(1).

Sunkaraneni VS, Jones SE. Topical anaesthetic or vasoconstrictor preparations for flexible fibre optic nasal pharyngoscopy and laryngoscopy. Cochrane Database of Systematic Reviews. 2011(3).

Therapists, Position paper. 2020

Todd S, Walsted ES, Grillo L, Livingston R, Menzies Gow A, Hull JH. Novel assessment tool to detect breathing pattern disorder in patients with refractory asthma. Respiriology. 2018 Mar;23(3):284-90.

Traister RS, Fajt ML, Landsittel D, Petrov AA. A novel scoring system to distinguish vocal cord dysfunction from asthma. The Journal of Allergy and Clinical Immunology: In Practice. 2014 Jan 1;2(1):65-9.

Van Doorn P, Folgering H, Colla P. Control of the end-tidal PCO₂ in the hyperventilation syndrome: effects of biofeedback and breathing instructions compared. Bulletin europeen de physiopathologie respiratoire. 1982 Nov 1;18(6):829-36.

Vertigan AE, Bone SL, Gibson PG. Development and validation of the Newcastle laryngeal hypersensitivity questionnaire. Cough. 2014 Dec;10(1):1-3.

Vertigan AE, Bone SL, Gibson PG. Laryngeal sensory dysfunction in laryngeal hypersensitivity syndrome. Respiriology. 2013 Aug;18(6):948-56.

Vertigan AE, Theodoros DG, Gibson PG, Winkworth AL. Efficacy of speech pathology management for chronic cough: a randomised placebo controlled trial of treatment efficacy. *Thorax*. 2006 Dec 1;61(12):1065-9.

Wallace S, McLaughlin C, Clayton J, Coffey M, Ellis J, Haag R, Howard A, Marks

H, Zorko R. Fiberoptic Endoscopic evaluation of Swallowing (FEES): The role of speech and language therapy. London: Royal College of Speech and Language

Therapists, Position paper. 2020

Wallace S, McLaughlin C, Clayton J, Coffey M, Ellis J, Haag R, Howard A, Marks H, Zorko R. Fiberoptic Endoscopic Evaluation of Swallowing (FEES): The role of speech and language therapy. London: Royal College of Speech and Language Therapists, Competency framework and training log. 2020.

The Voice Symptom Scale (VoiSS) and the Vocal Handicap Index (VHI): a comparison of structure and content. *Clinical Otolaryngology and Allied Sciences*. 2004 Apr;29(2):169-74.

Witek TJ, Mahler DA. Minimal important difference of the transition dyspnoea index in a multinational clinical trial. *European Respiratory Journal*. 2003 Feb 1;21(2):267-72.

We would like to acknowledge that this guidance has been written on behalf of the Royal College of Speech and Language Therapists (RCSLT) by:

- Jemma Haines MBE, (lead author), consultant speech and language therapist and RCSLT national adviser, Manchester Airways Service, Northwest Lung Centre, Wythenshawe Hospital, Manchester University NHS Foundation Trust, Manchester
- Karen Esposito, highly specialist speech and language therapist, Sheffield Teaching Hospitals NHS Foundation Trust
- Jen Butler (nee Murphy), highly specialist speech and language therapist, Newcastle upon Tyne Hospitals NHS Foundation Trust
- Nicola Pargeter, principal speech and language therapist, University Hospitals Birmingham NHS Foundation Trust
- Julia Selby, consultant speech and language therapist, Upper Airways Service, Royal Brompton Hospital
- Claire Slinger, consultant speech and language therapist, Lancashire Chest Centre, Lancashire Teaching Hospitals NHS Foundation Trust
- Hannah Lever, senior specialist speech and language therapist, Lancashire Chest Centre, Lancashire Teaching Hospitals NHS Foundation Trust

Key organisations

- **British Thoracic Society**
- **Association of Respiratory Nurse Specialists**
- **Association of Chartered Physiotherapists in Respiratory Care**
- **British Laryngology Society**
- **ENT-UK**