

References

- Abrams, H (2017) Hearing Loss and Associated Comorbidities: What Do We Know? *The Hearing Review*. <http://www.hearingreview.com/2017/11/hearing-loss-associated-comorbidities-know/> [Accessed 4 December 2019].
- Ayres, J (2005) Sensory Integration and the child. Western Psychological Services.
- Fortnum, H and Davis, A (1997) Epidemiology of Permanent Childhood Hearing Impairment in Trent Region, 1985–1993. *British Journal of Audiology*. 31 (6), 409-446.
- Hindley, P A et al (1994) Psychiatric disorder in deaf and hearing impaired children and young people: a prevalence study. *Journal of Child Psychology and Psychiatry*. 35, 917-934.
- Lin F R et al (2011). Hearing loss and incident dementia. *Archives of Neurology*. 68 (2), 214-220.
- Macaulay C E, and Ford R M (2006) Language and theory-of-mind development in prelingually deafened children with cochlear implants: a preliminary investigation. *Cochlear Implants International*. 7(1):1-14
- McConkey Robbins, A (2009) Rehabilitation after cochlear implantation. U: J K Niparko (ur.). *Cochlear implants: principles and practices* (str. 269-312) (2nd ed). Philadelphia: Wolters Kluwer/ Lippincott Williams and Wilkins.
- NICE (2019) Cochlear implants for children and adults with severe to profound deafness. Available from: <https://www.nice.org.uk/guidance/ta566/chapter/2-Clinical-need-and-practice> [Accessed 26 August 2020].